



OPEN ACCESS

*CORRESPONDENCE

Brian A. Cobb,
✉ brian.cobb@case.edu
Paolo Cravedi,
✉ paolo.cravedi@amssm.edu

†These authors share first authorship

‡These authors share senior authorship

RECEIVED 25 February 2026
REVISED 01 May 2026
ACCEPTED 11 May 2026
PUBLISHED 12 June 2026

CITATION

Noble J, Glendenning LM, Dard C, Bourdin A, Crespo M, Maggiore U, Inwood AR, Carlson GC, Cobb BA and Cravedi P (2026) Not all antibodies are created equal: total IgG glycosylation and severity of antibody-mediated rejection in kidney transplantation.
Transpl. Int. 39:16461.
doi: 10.3389/ti.2026.16461

COPYRIGHT

© 2026 Noble, Glendenning, Dard, Bourdin, Crespo, Maggiore, Inwood, Carlson, Cobb and Cravedi. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](#). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Not all antibodies are created equal: total IgG glycosylation and severity of antibody-mediated rejection in kidney transplantation

Johan Noble ^{1,2,3†}, Leandre M. Glendenning ^{4†}, Celine Dard ⁵, Anne Bourdin ⁵, Marta Crespo ⁶, Umberto Maggiore ⁷, Ari R. Inwood ⁴, Grace C. Carlson ⁴, Brian A. Cobb ^{4*‡} and Paolo Cravedi ^{1*‡}


¹Translational Transplant Research Center (TTRC), Icahn School of Medicine at Mount Sinai, Precision Immunology Institute, New York, NY, United States, ²Nephrology, Hemodialysis, Apheresis and Kidney Transplantation Department, University Hospital Grenoble, Grenoble, France, ³Inserm U 1209, CNRS UMR 5309, Team Cell Dynamics, Immunity, Metabolism & Cancer, Institute for Advanced Biosciences, University Grenoble Alpes Grenoble, France, ⁴Department of Pathology, Case Western Reserve University School of Medicine, Cleveland, OH, United States, ⁵Etablissement Français du Sang, Grenoble-Alpes, France, ⁶Department of Nephrology, Hospital Ddel Mar, Nephropathies Research Group, Hospital del Mar Research Institute, Barcelona, Spain, ⁷Nephrology Unit, Department of Medicine and Surgery, University Hospital of Parma, Parma, Italy

Antibody-mediated rejection (AMR) is a leading cause of kidney transplant failure and is primarily driven by donor-specific anti-HLA antibodies (DSA), although DSA presence alone does not fully explain the heterogeneity of AMR severity. We prospectively studied 79 kidney transplant recipients from 2 European centers, including 29 with active AMR (aAMR), 29 with chronic-active AMR (caAMR), and 21 controls without rejection, to investigate the association between total Immunoglobulin-G (IgG) glycosylation profiles and AMR. IgG glycosylation was quantified using lectin-based ELISA, assessing relative levels of mannose, core fucose, α 2,6-linked sialic acid, and bisecting N-acetylglucosamine (GlcNAc). Bisecting GlcNAc levels were higher in caAMR compared with controls and aAMR (both $p < 0.001$), while core fucosylation and mannose levels were increased in aAMR and caAMR relative to controls. Higher levels of core fucose, mannose, and α 2,6-sialylation were associated with increasing glomerulitis severity. We found that bisecting GlcNAc (PHA-E/Fc) was significantly associated with both higher g- and cg-score (in ordinal models, OR = 2.1 [95% CI: 1.2–4.1; $p = 0.017$], and OR = 2.0 [95% CI: 1.1–4.3; $p = 0.046$], respectively). Mannose level (ConA/Fc) was significantly associated with higher g-score (2.1 [1.1–4.2], $p = 0.017$). These findings indicate that distinct total IgG glycosylation features are independently associated with specific histological patterns of AMR severity.

KEYWORDS

antibody-mediated rejection, glycosylation, immunoglobulin G, N-acetylglucosamine, renal transplant

Not all alloantibodies are created equal: IgG glycosylation and severity of antibody-mediated rejection in kidney transplantation



GRAPHICAL ABSTRACT

Noble, et al. *Transpl. Int.* 2026doi: [10.3389/ti.2025.16461](https://doi.org/10.3389/ti.2025.16461)

Introduction

Antibody-mediated rejection (AMR) is a major cause of kidney transplant loss [1]. Although the pathophysiology of AMR is not fully understood, donor-specific anti-HLA antibodies (DSA) are thought to play an important pathogenic role. When present, these antibodies target endothelial graft cells, leading to their direct activation [2], the initiation of the complement cascade, and the recruitment of immune cells [3, 4]. Antibodies can also promote endothelial injury through antibody-dependent cellular cytotoxicity (ADCC) and the release of pro-inflammatory cytokines, both events that are primarily mediated by natural killer (NK) cells [5–8].

While DSA are central to AMR pathogenesis - triggering complement-dependent cytotoxicity (CDC), antibody-dependent cellular cytotoxicity (ADCC), and proinflammatory injury - not all patients with DSA experience graft loss [9]. This disconnect suggests that titer of DSA alone fails to explain clinical outcomes.

Evidence has shown that post-translational modification of antibodies affects their function [10, 11]. The type and level of antibody glycosylation can influence its ability to activate complement and induce ADCC, as well as their stability and circulating half-life [12]. Glycosylation of Immunoglobulin G (IgG) involves the addition of carbohydrate at a conserved site in the Fc

region (Asn297), and, less frequently, in the Fab region [13]. Unlike the conserved site in the Fc region, glycan sites within the Fab region are primarily introduced during somatic hypermutation. N-linked glycosylation starts with the transfer of a high mannose glycan from a dolichol-glycan donor in the endoplasmic reticulum (ER) and is critical for protein folding quality control (Supplementary Figure 1A). Upon exit from the ER and entry into the Golgi apparatus, the high mannose structures are matured by sequential removal of all but three mannoses, leaving a pentasaccharide core (Supplementary Figure 1B), and then rebuilt through the addition of N-acetylglucosamine (GlcNAc) residues (Supplementary Figure 1C) and varying degrees of additional capping glycans (Supplementary Figure 1D). This results in complex N-glycans carrying varied amounts of GlcNAc branching, core fucosylation, galactose and sialic acid residues. Glycan structures found on IgG are typically bi-antennary, and may contain bisecting GlcNAc (Supplementary Figure 1D), a branching modification that influences IgG function and enhances Fc receptor binding. Functionally, sialylation (addition of terminal N-acetylneuraminic acid with an α 2,6 linkage to the underlying galactose) and core fucosylation (fucose attached with an α 1,6 linkage to the asparagine-proximal GlcNAc) are associated with reduced ADCC properties by reducing Fc gamma receptors (Fc γ R)III affinity [14].

IgG glycosylation profile has been associated with changes in the severity of various autoimmune diseases [15]. A study in kidney transplant recipients showed that low sialylation and high bisecting GlcNAc of anti-donor specific HLA antibodies (DSA) were associated with higher risk of AMR [16]. However, this study assessed only post-translational modifications in IgG3. Herein, we hypothesized that total IgG glycosylation profile may be associated with the risk and the severity of both AMR and caAMR.

Abbreviations: AMR, Antibody-mediated rejection; ADCC, Antibody-dependent cellular cytotoxicity; cg, glomerular chronicity score; ConA, Concanavalin A; DSA, Donor-specific Antibody; g, glomerulitis; GlcNAc, N-acetylglucosamine; IgG, Immunoglobulin G; LCA, Lens culinaris agglutinin; ptc, peritubular capillaritis; MVI, microvascular inflammation; Neu5Ac, sialic acid; SNA, Sambucus nigra agglutinin.

Materials and methods

Study population

We used serum samples from prospectively followed-up kidney transplant recipients from two centers (Grenoble-Alpes, Grenoble, France and Hospital Del Mar of Barcelona, Barcelona, Spain) between January 2016 and May 2024 with a biopsy-proven AMR and control kidney transplant patients with no signs of microvascular inflammation (MVI). All samples were collected at the time of biopsy. All control biopsies were per protocol, taken at 3 months, 1, and 3 years post-transplant. All biopsies were scored according to Banff 2022 and the indications were protocol biopsies or for cause [17]. In brief, AMR was defined by the presence of MVI, i.e., glomerulitis (g) and peritubular capillaritis (ptc). If the MVI score was 2 or above, AMR was confirmed by the presence of C4d deposition in peritubular capillaries and vasa recta and/or by the presence of HLA DSA. If other features of AMR, such as acute thrombotic microangiopathy were present, but the MVI score was below 2, AMR was confirmed by the presence of C4d deposition, regardless of DSA status. C4d deposition was quantified using anti-C4d antibody on formalin-fixed, frozen paraffin-embedded sections. Chronic active AMR (caAMR) was defined by the presence of active lesions (including C4d) and chronic lesions (transplant glomerulopathy (cg) and/or peritubular capillary multilayering).

All patients signed an informed consent form. For Grenoble Hospital, all medical data were collected from the Grain-DB database [CNIL (French National committee for data protection) approval number 1987785v0 for the Nephrology and Kidney transplantation department of Grenoble-France]. The study was also approved by the Institutional Review Board of the Consorci Mar Parc de Salut de Barcelona (Research Project No. 2020/9117/I) and was conducted in accordance with the principles of the Declaration of Helsinki and the Declaration of Istanbul.

Anti-HLA antibody measurement

Anti-HLA antibodies were measured at the time of biopsy in all patients using One Lambda-ThermoFisher Labscreen, Class I, and Class II or Lifecodes LSA Class I and/or Class II test (Immucor GTI Diagnostics, Inc[®]). Antibody specificity was identified using a Luminex[®] 200 Instrument (Luminex Corporation, Austin, TX). The MFI threshold of positivity for a DSA was 500 for patients followed-up in Grenoble and 750 for those Barcelona. The sera used for the post-translational analysis were leftovers from the anti-HLA analysis at the Etablissement Français du Sang (EFS) of Grenoble, HLA department and study-dedicated sera for patients enrolled in Barcelona.

Immunoglobulin G purification

We purified total serum IgG using Protein A IgG Purification Kit (Pierce[™], Thermo Scientific[™], CAT: 44667) and following manufacturer's instructions. Briefly, the procedure consisted of passing the sera sample into the column and then eluting the IgG bound to the resin using an elution buffer (pH 2.8). For each sample, IgG concentration was measured by absorbance at 280 nm and purity was confirmed using SDS-PAGE.

Quantification of glycosylation by lectin ELISA

Purified IgG were diluted to 1 ng/ μ L in PBS, plated into a 96-well high-binding ELISA plate (Microton High Binding; Greiner BioOne) and incubated overnight at 4 °C. The plate was blocked with carbohydrate free blocking solution (Vector Labs) for 1 h at room temperature. Biotinylated lectins (Vector Labs) were diluted to 1 μ g/mL in carbohydrate-free blocking solution and incubated for 1 h at room temperature. Different biotinylated lectins were used: Sambucus nigra agglutinin (SNA): binds to α 2,6-linked Neu5Ac and is used to quantify the sialylation of IgG. Concanavalin A (ConA): binds to high mannose and hybrid N-glycans and correlates with more immature IgG. Lens culinaris agglutinin (LCA): binds to core fucosylated N-glycans. Phaseolus vulgaris Erythroagglutinin (PHA-E) binds to bisecting GlcNAc to quantify bisecting GlcNAc on IgG. Anti-Fc measures the amount of immobilized IgG to normalize the results. The amount of IgG coated was 0, 100ng and 200 ng to assess saturation of the signal, and we thereafter used the results of 100 ng for consistency. Signal was detected using europium-conjugated streptavidin (Perkin Elmer) and time-resolved fluorescence as measured in a Victor Nivo plate reader.

Statistical analysis

Normally distributed quantitative variables were expressed as mean \pm standard deviation (SD). Non-normally distributed quantitative variables were expressed as median [interquartile ranges (IQR)]. All glycosylation values were non-normally distributed (p-value from Shapiro-Wilk Test <0.005). Qualitative variables were expressed as numbers and percentages. For statistical comparison between multiple groups, we used Kruskal-Wallis test. For statistical comparisons between two groups, we used unpaired Mann-Whitney U Tests. The chi-squared test was used to compare categorical data. Statistical significance was defined as a two-sided p-value <0.05.

To identify factors associated with each ordinal outcome variable (i.e., scores of Banff components), we employed ordinal logistic regressions using the proportional odds model using the *clm* function from the ordinal R package (R version 4.4.2, <https://www.r-project.org>). Ordinal logistic regression models for g- cg- and pc-scores were fitted for each glycan, and adjusted for DSA, recipient age, sex, delay of biopsy since transplantation, and eGFR at biopsy whenever they were statistically significant to avoid overfitting given the limited sample size. These covariates were included, as they may affect both glycosylation and histological outcomes independently of any direct glycosylation-mediated effect [18, 19]. To visualize the ordinal regression model for cg score, we plotted the fitted predicted probability for each cg score level as a function of standard unit change of bisecting GlcNAc (PHA-E/Fc).

Results

Baseline characteristics of the study population

The study included 79 kidney transplant recipients, 29 (36.7%) with biopsy-proven aAMR, 29 (36.7%) with caAMR, and 21 (26.6%)

TABLE 1 Baseline characteristics of study participants.

Characteristics	CTL (N = 21)	aAMR (N = 29)	caAMR (N = 29)	p-value
Age - years	51.7 ± 12	43.7 ± 17	43.4 ± 17	0.179
Gender M/F ratio	3.2	0.9	1.7	0.125
History of previous transplantation - N (%)	0	8 (27.6)	3 (10.3)	0.065
Nephropathy - N (%)				0.522
Diabetes	1 (4.7)	0	2 (6.9)	
Drug-related	1 (4.7)	0	0	
Nephrotic disease	1 (4.7)	3 (10.3)	2 (6.9)	
Genetic	1 (4.7)	1 (3.4)	0	
Uropathy	1 (4.7)	2 (6.9)	3 (10.3)	
Unknown	7 (33.2)	7 (24.1)	5 (17.24)	
Glomerulonephritis	1 (4.7)	7 (24.1)	6 (20.7)	
TMA	1 (4.7)	1 (3.4)	1 (3.4)	
Hypertensive nephropathy	2 (9.5)	3 (10.3)	3 (10.3)	
Interstitial nephropathy	2 (9.5)	1 (3.4)	2 (6.9)	
Acute tubular necrosis	1 (4.7)	0	0	
Polycystic disease	4 (19)	2 (6.9)	4 (13.8)	
Time of biopsy - months after transplant	38.5 [3–43]	10.9 [2–58]	47.3 [35–149]	0.004
PRA (%)	11 ± 28	49 ± 44	30 ± 45	0.021
Living donor - N (%)	4 (19)	7 (24.1)	6 (6.9)	0.902
Donor age - years	55.2 ± 14	49.5 ± 16	45.8 ± 16	0.192
HLA mismatch A/B/DQ - number	5.0 [3–6]	6.0 [5–7]	5.5 [5–7]	0.218
History of diabetes - N (%)	1 (4.7)	0	2 (6.9)	0.329
Anti-thymoglobulin induction - N (%)	9 (42.7)	21 (72.4)	20 (68.9)	0.725
MVI ≥2 - N (%)	0	24 (82.7)	28 (96.5)	<0.001
C4d positivity- N (%)	0	22 (75.8)	10 (34.5)	<0.001
Donor-specific antibodies - N (%)	1 (4.7)	23 (79.3)	21 (72.4)	<0.001
cg-score positivity- N (%)	0	0	29 (100)	<0.001
eGFR at the time of biopsy - mg/dL	43 [31–63]	45 [32–66]	36 [23–49]	0.145
Proteinuria at rejection - g/g	0.1 [0.1–0.4]	0.3 [0.1–0.7]	0.8 [0.3–2.0]	<0.001

FSGS, Focal segmental glomerulosclerosis; HLA, Human Leukocyte Antigen; IgAN, IgA nephropathy; MVI, microvascular inflammation; PRA, panel reactivity antibody; eGFR, estimated glomerular filtration rate; TMA, thrombotic microangiopathy.


controls with no histological signs of rejection at per-protocol biopsy (Table 1). There was no difference among patients with aAMR, caAMR and controls in the main characteristics such as the age at the time of biopsy, gender, previous transplantation number, type of donor, HLA mismatches, history of diabetes, and induction therapy. The median time between the transplantation and the biopsy was 0.9 years [0.1–4.8] in the aAMR group, 3.9 [2.9–12.3] for the caAMR group, and 3.2 [0.3–3.6] for the control group, $p = 0.004$. The immunosuppressive regimen consisted of tacrolimus, mycophenolate mofetil, and steroids for all patients. Patients with aAMR or caAMR had higher proteinuria than controls (Table 1).

Twenty-three aAMR patients (79.3%) had donor-specific antibody (DSA) at the time of the biopsy, compared to 21 (72.4%) in the caAMR group ($p = 0.539$). One patient in the control group had DSA. The number of DSA was also similar between the two rejection groups (1.5 ± 0.9 vs. 1.4 ± 0.7 , $p =$

0.539) and most of DSA were class II (81% versus 85%, $p = 0.731$) for aAMR and caAMR, respectively. The mean fluorescence intensity (MFI) of the immunodominant DSA was also similar (8226 ± 6549 vs. 4915 ± 5968 , $p = 0.192$).

Associations between IgG glycosylation and the risk of AMR

The quantification of IgG glycan composition (relative amounts of mannose, fucose, sialic acid, and bisecting GlcNAc) was assessed using the ConA, LCA, SNA, and PHA-E lectins, respectively, and normalized with the Fc levels (Figure 1). We found a positive correlation between mannose, core fucose, $\alpha 2,6$ -linked sialic acids. In contrast, bisecting GlcNAc was not significantly associated with the other glycoforms (Supplementary Figure 2). There was no



correlation between the glycan level and the total amount of IgG (data not shown).

We first compared the different IgG glycoforms among the 3 groups. The amounts of bisecting GlcNAc (PHA-E/Fc), core fucose (LCA/Fc), and mannose (ConA/Fc) were significantly lower in the control group than in the caAMR group. Bisecting GlcNAc (PHA-E/Fc) was significantly higher in the caAMR and aAMR groups compared to controls, and significantly higher in caAMR compared to aAMR patients (Figure 2A). Core fucose (LCA/Fc) was significantly higher in both caAMR and aAMR groups compared to controls (Figure 2B). Mannose (ConA/Fc) was significantly higher in caAMR than in controls (Figure 2C). There was no statistical difference among the 3 groups in the α 2,6-sialylation (SNA/Fc) levels (Figure 2D). When gathering aAMR and caAMR, we found higher level of bisecting GlcNAc ($p < 0.001$), of fucosylation ($p < 0.001$), and of Mannose ($p < 0.001$) in rejection patients compared to controls while sialylation levels were similar ($p = 0.41$).

Associations between IgG glycosylation and the severity of rejection

We then compared the amount of each glycoform with the severity of AMR using each of the AMR parameters listed in the Banff score (0–3), i.e., g-, cg-, and ptc- scores and C4d deposition intensity (0–3). We used cg-score to quantify the chronicity of lesions. We found no association between bisecting GlcNAc (PHA-E/Fc) and g-score increasing severity (Figure 3A). Core fucose (LCA/Fc), mannose (ConA/Fc) and α 2,6-sialic acids (SNA/Fc) levels in IgG were increased in samples from patients with the highest g-scores compared to those with lower scores (Figures 3B–D). Within AMR cases, core fucosylation was increasing with the severity of glomerulitis (Figure 3B). We also found that patients

with highest cg-score had increased bisecting GlcNAc (PHA-E/Fc), core fucose (LCA/Fc), mannose (ConA/Fc), and sialylation (SNA/Fc; Supplementary Figure 3) levels. Finally, there was a correlation between ptc-score severity and IgG glycan levels (Supplementary Figure 4).


Association between IgG glycosylation and DSA

Not all patients in our cohorts had DSA. Therefore, we tested the association between IgG glycosylation and DSA. Among the different glycosylation tested, we found no significant correlation between total IgG glycosylation and the presence of DSA in patients with AMR (Supplementary Figure 5A–D). In particular, the level of bisecting GlcNAc was not different between aAMR and caAMR patients with or without DSA (Supplementary Figure 5A).

We repeated the analyses comparing glycoform levels in controls, aAMR, and caAMR cohorts, by stratifying patients based on the presence of DSA. In DSA negative patients, caAMR remained significantly higher than controls for bisecting GlcNAc, total mannose and core fucose while aAMR remained significantly associated with total mannose and core fucose (Figures 4A–C). In DSA positive patients, bisecting GlcNAc was higher in caAMR compared to aAMR patients (Figure 4A). α 2,6-sialylation was not significantly different between controls and AMR conditions, regardless of DSA status (Figure 4D).

Multivariable analysis of factors associated with histologic features of AMR

Finally, we tested whether the IgG glycosylation and AMR severity are independently associated.




We found that bisecting GlcNAc (PHA-E/Fc) was significantly associated with both higher g- and gc-scores (in ordinal models, OR = 2.1 [95% CI: 1.2–4.1; $p = 0.017$], and OR = 2.0 [95% CI: 1.1–4.3; $p = 0.046$], respectively). Mannose level (ConA/Fc) was significantly associated with higher g-score (2.1 [1.1–4.2], $p = 0.017$) (Table 2). Core fucosylation and sialylation were not significantly associated with AMR severity in multivariable analyses. In Table 2 DSA might seem protective with respect to cg-score (OR less than 1), but this multivariable finding is artifactual as shown by the visual inspection of the predicted probability from the ordinal regression model (Figure 5). In DSA-negative patients, high bisecting GlcNAc predicted severe glomerulopathy (cg = 3), while low levels were protective (cg = 0). This dose-response relationship was attenuated in DSA-positive patients, where high bisecting GlcNAc barely influenced glomerulopathy severity. The apparent protective effect of DSA in multivariable models is therefore artifactual, likely reflecting DSA's dominant role that masks glycosylation-mediated injury. Bisecting GlcNAc and

DSA appear to represent independent but competing pathogenic mechanisms.

Discussion

In this study, we found an association between total IgG core-fucosylation, bisecting GlcNAc, and detectable mannose, and the presence of biopsy-proven AMR in kidney transplant recipients. In contrast, IgG α 2,6-sialylation was not associated with AMR in our cohort. Moreover, bisecting GlcNAc was significantly associated with both glomerulitis (g-score) and chronic glomerulopathy (cg-score). Interestingly, this association appeared independent of DSA status and may represent a competing pathogenic mechanism.

An inverse association between DSA presence and cg-score emerged in the multivariable model. Exploratory interaction analyses suggest that this finding likely reflects effect-modification




rather than a protective role of DSA. The association between higher bisecting GlcNAc levels and severe chronic glomerulopathy was mainly observed in DSA-negative patients, whereas in DSA-positive individuals, glycosylation levels did not substantially alter the predicted probability of severe lesions. These findings suggest that the association between IgG glycosylation and chronic glomerulopathy may be primarily observable in the absence of detectable DSA, whereas in DSA-positive patients, the pathogenic signal associated with DSA may dominate and attenuate the relative contribution of glycosylation levels.

Our findings confirm and expand results from prior smaller studies. Pernin et al. performed mass spectrometry on isolated IgG3 *de novo* DSA and showed that higher bisecting GlcNAc glycosylation was higher in AMR patients overall compared to patients DSA+, but without histological AMR [16]. If mass spectrometry allows a precise characterization of glycoforms, it can also exhibit higher technical variability compared to other methods, potentially affecting reproducibility [20]. ELISA, on the contrary, is well-suited for large-scale studies due to its simplicity and ability to process many samples simultaneously, is generally less expensive and requires less specialized equipment compared to mass spectrometry, and can be standardized across different laboratories, enhancing reproducibility [21].

Our findings are in line with the literature about bisecting GlcNAc. Bisecting GlcNAcylation levels can modify the Fc activity of IgG by influencing its binding affinity to Fc γ R_s. Specifically, the presence of a bisecting GlcNAc moiety has been shown to inhibit Fc core-fucosylation and enhance the binding of Fc to Fc γ RIIIa, an activating Fc γ receptor [22, 23]. This enhancement in binding affinity can lead to increased ADCC activity, which conceivably results in increased glomerular injury induced by HLA and non-HLA antibodies.

In the study by Pernin et al., sialylation of IgG3 DSA was found to be associated with a reduced risk of AMR. Sialylation of total IgG assessed at the time of transplantation was also associated with reduced AMR risk during the follow-up in another study [24]. In our cohort, we did not find any correlation between IgG sialylation and the presence of AMR. Together, these data suggest a reduced activity of HLA and non-HLA antibodies in kidney transplantation when associated with a higher sialylation. This hypothesis is supported by the literature, as sialylation of the IgG Fc domain has been shown to impair complement-dependent cytotoxicity (CDC) by reducing the binding of C1q [25], decreasing ADCC (in the presence of core fucosylation) by reducing the binding affinity of IgG to Fc γ RIIIa [26], and exhibiting anti-inflammatory effects [27].




The significance of our data extends to individuals with autoimmune diseases, where alterations in IgG glycosylation have been well-documented. For instance, in rheumatoid arthritis, there is a notable decrease in galactosylation and sialylation, and increased fucosylation in IgG during active phases of the disease [28], suggesting that these glycosylation changes can serve as biomarkers for disease activity and therapeutic response. Similarly, systemic lupus erythematosus patients exhibit decreased IgG galactosylation and sialylation, increased bisecting GlcNAc, and decreased core fucose [29, 30]. These glycosylation changes are associated with disease activity and specific clinical manifestations, such as nephritis and the onset of antinuclear antibodies. IgG glycosylation patterns, including sialylation, bisecting GlcNAcylation, fucosylation, and mannosylation, are significantly altered in various other autoimmune and inflammatory diseases, and these changes have been proposed as potential biomarkers for disease activity and severity [31, 32].

One limitation of our study is that we analyzed total IgG glycosylation, instead of DSA-specific IgG glycosylation, as they may theoretically differ. Of note, one single patient in the control group had DSA, further limiting the power of our analyses to infer the impact of total IgG glycosylation, DSA and AMR severity. However, Sonneveld et al. demonstrated that the Fc-glycosylation profiles of total serum IgG1 and IgG3 were highly similar to those of antigen-specific anti-K antibodies [33]. In that study, the authors analyzed the Fc-tail glycosylation of both total serum and antigen-specific IgG1 and IgG3 using mass spectrometry and found that the glycosylation patterns were consistent across both types, with little individual variability. Using an integrated method for simultaneous protein quantitation and glycosylation profiling of antigen-specific IgG1, Gijze et al. also found that the glycosylation profiles of antigen-specific IgG1 are comparable to those of total IgG1 in a large coronavirus vaccination cohort [21]. Of note, we found that the differences in bisecting

TABLE 2 Multivariable ordinal regression analysis of the association between clinical variable, IgG post-translational modifications, and AMR severity scores in the overall cohort.

Variables	g-score		ptc-score		cg-score	
	OR (95% CI)	p-value	OR (95% CI)	p-value	OR (95% CI)	p-value
Presence of DSA						
Bisecting GlcNAc	2.1 (1.2–4.1)	0.017			2.0 (1.1–4.3)	0.046
Core fucose						
α2,6 sialic acid						
Mannose	.1 (1.1–4.2)	0.017				
Recipient age (years)						
Recipient gender (male)			4.1 (1.2–15.7)	0.027	3.7 (1.1–13.8) ^{S,&,E,§}	0.038
eGFR (ml/min/1.73m ²)*					0.9 (0.9–0.9) ^{S,&,E}	0.02
Delay post-transplant (year)					1.1 (1.0–1.2) ^{S,&,E}	0.009
DSA						

Bisecting GlcNAc (PHA-E/Fc ratio); Core fucose (LCA/Fc ratio); α2,6 sialic acid (SNA/Fc ratio); mannose (ConA/Fc ratio), DSA: Donor specific Antibody; eGFR: estimated Glomerular Filtration Rate. *At rejection. ^SModel with bisecting GlcNAc. ^EModel with Core fucosylation. [&]Model with Sialylation. [§]Model with Mannose.



GlcNAcylation persist in patients with and in those without DSA, suggesting that although DSA may have a unique glycosylation profile, some key immunomodulatory glycoforms are probably shared across antibody specificities. However, *ad hoc* studies are needed to formally test this working model. Glycosylation of total IgG may capture information from non-HLA anti-donor antibodies that are hard to capture in routine clinical testing, or it may reflect a more systemic state of inflammation. Further studies are necessary to confirm and investigate these hypotheses.

In conclusion, we found a significant, independent correlation between total IgG glycosylation and AMR in kidney transplant recipients. More specifically, we showed the association of IgG bisecting GlcNAc regardless of the presence of DSA, with glomerulitis and chronic glomerulopathy in graft biopsy. Therefore, our findings provide the rationale for larger studies testing the hypothesis that total IgG glycosylation can be used as a biomarker for the risk of aAMR and caAMR. Strategies to target IgG glycosylation may also be leveraged therapeutically to improve the outcomes of transplant recipients.

Data availability statement

The original contributions presented in the study are included in the article/supplementary material, further inquiries can be directed to the corresponding authors.

Ethics statement

The studies involving humans were approved by CNIL approval number 1987785v0 and Institutional Review of Barcelona, project 2020/9117/l. The studies were conducted in accordance with the local legislation and institutional requirements. The participants provided their written informed consent to participate in this study.

Author contributions

PC and BC designed the study, JN, LG, AI, and GC performed the experiments. JN wrote the manuscript. CD and AB collected the samples. UM performed statistical analysis. All authors contributed to the article and approved the submitted version.

Funding

The author(s) declared that financial support was received for this work and/or its publication. This work was funded by National Institutes of Health grants to BAC (GM115234), LMG (AI178953), GCC (AI089474). MC had funding from Instituto de Salud Carlos III with FIS-FEDER (PI23/00061 and BA24/00014) and RICORS program RD24/0004/0003, both co-financed by the European Union.

References

- Lefaucheur C, Loupy A. Antibody-mediated rejection of solid-organ allografts. *N Engl J Med* (2018) 379(26):2580–2. doi:10.1056/NEJMc1813976
- Valenzuela NM, Mulder A, Reed EF. HLA class I antibodies trigger increased adherence of monocytes to endothelial cells by eliciting an increase in endothelial P-selectin and, depending on subclass, by engaging FcγRs. *J Immunol* (2013) 190(12):6635–50. doi:10.4049/jimmunol.1201434
- Bailly E, Macedo C, Ossart J, Louis K, Gu X, Ramaswami B, et al. Interleukin-21 promotes Type-1 activation and cytotoxicity of CD56dimCD16bright natural killer cells during kidney allograft antibody-mediated rejection showing a new link between adaptive and innate humoral allo-immunity. *Kidney Int* (2023) 104(4):707–23. doi:10.1016/j.kint.2023.04.024
- Nevarez-Mejia J, Jin YP, Pickering H, Parmar R, Valenzuela NM, Sosa RA, et al. Human leukocyte antigen class I antibody-activated endothelium promotes CD206+ M2 macrophage polarization and MMP9 secretion through TLR4 signaling and P-selectin in a model of antibody-mediated rejection and allograft vasculopathy. *Am J Transpl* (2024) 24(3):406–18. doi:10.1016/j.ajt.2023.10.020
- Charreau B. Cellular and molecular crosstalk of graft endothelial cells during AMR: effector functions and mechanisms. *Transplantation* (2021) 105(11):e156–67. doi:10.1097/TP.00000000000003741
- Koenig A, Mezaache S, Callemeyn J, Barba T, Mathias V, Sicard A, et al. Missing self-induced activation of NK cells combines with non-complement-fixing donor-specific antibodies to accelerate kidney transplant loss in chronic antibody-mediated rejection. *J Am Soc Nephrol* (2021) 32(2):479–94. doi:10.1681/ASN.2020040433
- Chambon M, Koenig A. NK cells: not just followers but also initiators of chronic vascular rejection. *Transpl Int* (2024) 37:13318. doi:10.3389/ti.2024.13318
- Charmetant X, Pettigrew GJ, Thauan O. Allorecognition unveiled: integrating recent breakthroughs into the current paradigm. *Transpl Int* (2024) 37:13523. doi:10.3389/ti.2024.13523
- Koenig A, Chen CC, Marçais A, Barba T, Mathias V, Sicard A, et al. Missing self triggers NK cell-mediated chronic vascular rejection of solid organ transplants. *Nat Commun* (2019) 10(1):5350. doi:10.1038/s41467-019-13113-5
- van de Bovenkamp FS, Hafkenscheid L, Rispens T, Rombouts Y. The emerging importance of IgG Fab glycosylation in immunity. *J Immunol* (2016) 196(4):1435–41. doi:10.4049/jimmunol.1502136
- Krištić J, Lauc G. The importance of IgG glycosylation-what did we learn after analyzing over 100,000 individuals. *Immunol Rev* (2024) 328(1):143–70. doi:10.1111/imr.13407
- Quast I, Peschke B, Lünemann JD. Regulation of antibody effector functions through IgG Fc N-glycosylation. *Cell Mol Life Sci* (2017) 74(5):837–47. doi:10.1007/s00018-016-2366-z
- Cobb BA. The history of IgG glycosylation and where we are now. *Glycobiology* (2020) 30(4):202–13. doi:10.1093/glycob/cwz065
- Golay J, Andrea AE, Cattaneo I. Role of Fc core fucosylation in the effector function of IgG1 antibodies. *Front Immunol* (2022) 13:929895. doi:10.3389/fimmu.2022.929895
- Plomp R, Ruhaak LR, Uh HW, Reiding KR, Selman M, Houwing-Duistermaat JJ, et al. Subclass-specific IgG glycosylation is associated with markers of inflammation and metabolic health. *Sci Rep* (2017) 7(1):12325. doi:10.1038/s41598-017-12495-0

Acknowledgements

We thank Shivani Wadnerkar and Friederike Selbach for helping isolate IgG from patients' sera. A previous version of this manuscript has been deposited as a preprint on bioRxiv (<https://www.biorxiv.org/content/10.1101/2025.05.28.656600v1>).

Conflict of interest

The author(s) declared that this work was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Generative AI statement

The author(s) declared that generative AI was not used in the creation of this manuscript.

Any alternative text (alt text) provided alongside figures in this article has been generated by Frontiers with the support of artificial intelligence and reasonable efforts have been made to ensure accuracy, including review by the authors wherever possible. If you identify any issues, please contact us.

Supplementary material

The Supplementary Material for this article can be found online at: <https://www.frontierspartnerships.org/articles/10.3389/ti.2026.16461/full#supplementary-material>

16. Pernin V, Bec N, Beyze A, Bourgeois A, Szwarz I, Champion C, et al. IgG3 donor-specific antibodies with a proinflammatory glycosylation profile may be associated with the risk of antibody-mediated rejection after kidney transplantation. *Am J Transpl* (2022) 22(3):865–75. doi:10.1111/ajt.16904
17. Naesens M, Roufosse C, Haas M, Lefaucheur C, Mannon RB, Adam BA, et al. The Banff 2022 kidney meeting report: reappraisal of microvascular inflammation and the role of biopsy-based transcript diagnostics. *Am J Transpl* (2024) 24(3):338–49. doi:10.1016/j.ajt.2023.10.016
18. Wu Y, Zhang Z, Chen L, Sun S. Immunoglobulin G glycosylation and its alterations in aging-related diseases. *Acta Biochim Biophys Sin (Shanghai)* (2024) 56(8):1221–33. doi:10.3724/abbs.2024137
19. Baković MP, Selman MHJ, Hoffmann M, Rudan I, Campbell H, Deelder AM, et al. High-throughput IgG Fc N-glycosylation profiling by mass spectrometry of glycopeptides. *J Proteome Res* (2013) 12(2):821–31. doi:10.1021/pr300887z
20. Reiding KR, Bondt A, Hennig R, Gardner RA, O’Flaherty R, Trbojević-Akmačić I, et al. High-throughput serum N-Glycomics: Method comparison and application to study rheumatoid arthritis and pregnancy-associated changes. *Mol Cell Proteomics* (2019) 18(1):3–15. doi:10.1074/mcp.RA117.000454
21. Gijze S, Wasynczuk A, van Leeuwen L, Grobden M, van Gils MJ, Nouta J, et al. Simultaneous protein quantitation and glycosylation profiling of antigen-specific immunoglobulin G1 in large clinical studies. *J Proteome Res* (2024) 23(12):5600–5. doi:10.1021/acs.jproteome.4c00538
22. Zou G, Ochiai H, Huang W, Yang Q, Li C, Wang LX. Chemoenzymatic synthesis and Fcγ receptor binding of homogeneous glycoforms of antibody Fc domain. Presence of a bisecting sugar moiety enhances the affinity of Fc to FcγIIIa receptor. *J Am Chem Soc* (2011) 133(46):18975–91. doi:10.1021/ja208390n
23. Bosman GP, Stoof ID, Bastiaansen HP, Quarles van Ufford L, Dobruchowska JM, Langenbach JWH, et al. Bisecting N-Acetylglucosamine of the N-Glycan of immunoglobulin G does not affect binding to Fc gamma receptors. *ACS Chem Biol* (2025) 20(3):680–9. doi:10.1021/acscchembio.4c00807
24. Malard-Castagnet S, Dugast E, Degauque N, Pallier A, Soullillou JP, Cesbron A, et al. Sialylation of antibodies in kidney recipients with *de novo* donor specific antibody, with or without antibody mediated rejection. *Hum Immunol* (2016) 77(11):1076–83. doi:10.1016/j.humimm.2015.10.021
25. Quast I, Keller CW, Maurer MA, Giddens JP, Tackenberg B, Wang LX, et al. Sialylation of IgG Fc domain impairs complement-dependent cytotoxicity. *J Clin Invest* (2015) 125(11):4160–70. doi:10.1172/JCI82695
26. Li T, DiLillo DJ, Bournazos S, Giddens JP, Ravetch JV, Wang LX. Modulating IgG effector function by Fc glycan engineering. *Proc Natl Acad Sci U S A* (2017) 114(13):3485–90. doi:10.1073/pnas.1702173114
27. Ohmi Y, Ise W, Harazono A, Takakura D, Fukuyama H, Baba Y, et al. Sialylation converts arthritogenic IgG into inhibitors of collagen-induced arthritis. *Nat Commun* (2016) 7:11205. doi:10.1038/ncomms11205
28. Su Z, Xie Q, Wang Y, Li Y. Abberant immunoglobulin G glycosylation in rheumatoid arthritis by LTQ-ESI-MS. *Int J Mol Sci* (2020) 21(6):2045. doi:10.3390/ijms21062045
29. Vučković F, Krištić J, Gudelj I, Teruel M, Keser T, Pezer M, et al. Association of systemic lupus erythematosus with decreased immunosuppressive potential of the IgG glycome. *Arthritis Rheumatol* (2015) 67(11):2978–89. doi:10.1002/art.39273
30. Ramos-Martínez I, Ramos-Martínez E, Cerbón M, Pérez-Torres A, Pérez-Campos Mayoral L, Hernández-Huerta MT, et al. The role of B cell and T cell glycosylation in systemic lupus erythematosus. *Int J Mol Sci* (2023) 24(1):863. doi:10.3390/ijms24010863
31. Šimurina M, de Haan N, Vučković F, Kennedy NA, Štambuk J, Falck D, et al. Glycosylation of immunoglobulin G associates with clinical features of inflammatory bowel diseases. *Gastroenterology* (2018) 154(5):1320–33.e10. doi:10.1053/j.gastro.2018.01.002
32. Ząbczyńska M, Link-Lenczowski P, Pocheć E. Glycosylation in autoimmune diseases. *Adv Exp Med Biol* (2021) 1325:205–18. doi:10.1007/978-3-030-70115-4_10
33. Sonneveld ME, Koeleman CAM, Plomp HR, Wuhler M, van der Schoot CE, Vidarsson G. Fc-Glycosylation in human IgG1 and IgG3 is similar for both total and anti-red-blood cell Anti-K antibodies. *Front Immunol* (2018) 9:129. doi:10.3389/fimmu.2018.00129