



OPEN ACCESS

*CORRESPONDENCE

Helen Opdam,
✉ helen.opdam@austin.org.au

RECEIVED 17 July 2026
ACCEPTED 21 July 2026
PUBLISHED 29 July 2026

CITATION

Opdam H, Pérez-Blanco A, Gardiner D, Hasz R, Jansen NE, Le Dorze M, Sandiumenge A, Testa G, Then S-N, Zanierato M, Domínguez-Gil B, Oniscu GC, Cillo U and Martin DE (2026) Correction: Establishing standards for controlled donation after circulatory determination of death in adults: the Bucharest international European Society for Organ Transplantation consensus. *Transpl. Int.* 39:17395. doi: 10.3389/ti.2026.17395

COPYRIGHT

© 2026 Opdam, Pérez-Blanco, Gardiner, Hasz, Jansen, Le Dorze, Sandiumenge, Testa, Then, Zanierato, Domínguez-Gil, Oniscu, Cillo and Martin. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Correction: Establishing standards for controlled donation after circulatory determination of death in adults: the Bucharest international European Society for Organ Transplantation consensus

Helen Opdam^{1,2*}, Alicia Pérez-Blanco³, Dale Gardiner⁴, Richard Hasz⁵, Nichon Esther Jansen⁶, Matthieu Le Dorze^{7,8}, Alberto Sandiumenge^{9,10}, Giuliano Testa¹¹, Shih-Ning Then¹², Marinella Zanierato¹³, Beatriz Domínguez-Gil³, Gabriel C. Oniscu¹⁴, Umberto Cillo¹⁵ and Dominique E. Martin¹⁶

¹Department of Intensive Care, Austin Hospital, Heidelberg, VIC, Australia, ²Australian Organ and Tissue Authority, Canberra, ACT, Australia, ³Organizacion Nacional de Trasplantes, Madrid, Spain, ⁴Organ and Tissue Donation and Transplantation, NHS Blood and Transplant, Bristol, United Kingdom, ⁵Gift of Life Donor Program, Philadelphia, PA, United States, ⁶Department of Policy and Research, Dutch Transplant Foundation, Leiden, Netherlands, ⁷Department of Anesthesia and Critical Care, AP-HP, Hôpital Lariboisière, Paris, France, ⁸Université Paris Cité, INSERM U942 MASCOT, Paris, France, ⁹Vall d'Hebron Research Institute, Barcelona, Spain, ¹⁰Hospital Universitari Vall d'Hebron, Barcelona, Spain, ¹¹Simmons Transplant Institute, Baylor University Medical Center, Dallas, TX, United States, ¹²Australian Centre for Health Law Research, Faculty of Business & Law, Queensland University of Technology, Brisbane, QLD, Australia, ¹³Department of Anesthesia and Critical Care, AOU Città della Salute e della Scienza di Torino, Turin, Italy, ¹⁴Division of Transplantation Surgery, CLINTEC, Karolinska Institutet, Stockholm, Sweden, ¹⁵Unità di Chirurgia Epatobiliare e Trapianto Epatico, Università degli Studi di Padova, Padova, Italy, ¹⁶School of Medicine, Deakin University, Geelong, VIC, Australia

KEYWORDS

consensus, DCD (donation after circulatory death), deceased donation, donation, transplantation

A Correction on

Establishing standards for controlled donation after circulatory determination of death in adults: the Bucharest international European Society for Organ Transplantation consensus

by Opdam H, Pérez-Blanco A, Gardiner D, Hasz R, Jansen NE, Le Dorze M, Sandiumenge A, Testa G, Then S-N, Zanierato M, Domínguez-Gil B, Oniscu GC, Cillo U and Martin DE (2026). *Transpl. Int.* 39:16284. doi: 10.3389/ti.2026.16284

There was a mistake in [Table 1](#) as published. Incorrect percentages are included in two sections of the table. In addition, throughout the table, the percentages are shown to two decimal places, whereas they should be to a single decimal place. The corrected [Table 1](#) appears below.

The original version of this article has been updated.

TABLE 1 Expert panel demographics.

Panelist characteristics	First-wave n = 37	Second-wave n = 35
Age		
18–30 years	0.0%	0.0%
31–40 years	10.8%	11.4%
41–50 years	24.3%	22.9%
51–60 years	35.1%	37.1%
>60 years	29.7%	28.6%
Gender		
Male	64.9%	62.9%
Female	35.1%	37.1%
Country of employment		
Argentina	5.4%	5.7%
Australia	10.8%	11.4%
Canada	5.4%	5.7%
China	5.4%	2.9%
France	8.1%	8.6%
Italy	10.8%	11.4%
Japan	2.7%	0.0%
Netherlands	8.1%	8.6%
New Zealand	2.7%	2.9%
South Africa	2.7%	2.9%
Spain	8.1%	8.6%
Sweden	2.7%	2.9%
United Kingdom	10.8%	11.4%
United States	13.5%	14.3%
Uruguay	2.7%	2.9%
Primary specialty		
Intensivist/critical care physician	40.0%	39.6%
Transplant or donation coordinator	18.0%	18.8%
Transplant surgeon	14.0%	12.5%
Anaesthesiologist	6.0%	6.3%
Ethicist or legal scholar	6.0%	6.3%
Others*	16.0%	16.7%
Primary patient population		
Adult medicine	67.6%	65.7%
Both paediatric and adult medicine	16.2%	17.1%
Not applicable to my role	16.2%	17.1%

(Continued)

TABLE 1 Continued

Panelist characteristics	First-wave n = 37	Second-wave n = 35
Paediatrics	0.0%	0.0%
Primary sector of clinical practice for previous 12 months		
Public healthcare	73.0%	71.4%
Private healthcare	0.0%	0.0%
Both	16.2%	17.1%
Not Applicable	10.8%	11.4%
Type of hospital in which work		
Academic hospital	59.5%	60.0%
Non-academic hospital	8.1%	5.7%
Other academic centres and institutions	16.2%	17.1%
Other	16.2%	17.1%
Duration of experience with deceased donation or transplantation		
≥10 years	86.5%	85.7%
5–9 years	10.8%	11.4%
<5 years	2.7%	2.9%
Author contributions to academic publications in deceased organ donation or transplantation		
Yes	94.6%	94.3%
No	2.7%	2.9%
Not applicable	2.7%	2.9%
Participation as a principal investigator in a clinical trial relating to deceased organ donation or transplantation		
No	59.5%	62.9%
Yes	37.8%	34.3%
Not applicable to my role	2.7%	2.9%

*Values may not add up to 100% due to rounding errors.